

NAME _____

DATE _____

Module 5 Solving Linear Inequalities of One Variable
Lesson 7 Solving Problems Using Inequalities of One Variable



**independent
practice**

Solve each problem.

- A number, x , decreased by 5 is no more than 0. What are the possible values of the number?

- Jeb must spend at least \$97.50 on identical awards for 15 community volunteers. What is the least he can spend on each award?

- Each of seven members of the 4-H club raised equal amounts of money to attend the county fair. In all, the members raised over \$630. How much did each individual member raise?

- Dot Common, a computer tutoring service, charges \$10 to come to a customer's home and \$14 per hour to show customers how to use their computer. If a customer must spend less than \$55.50 on a single visit from Dot Common, how many hours of help can she receive?

- Giselle pays 30% of her weekly salary in taxes. She earns more than \$385 each month after taxes are taken from her paycheck. What is the minimum amount Giselle earns in a week **before** taxes are taken away?

- Many merchants require a minimum value for purchases made with credit cards because of the fees stores pay to credit card companies. If the items in a store are marked up 45% and the store requires a minimum of a \$5.00 for credit card purchases, what is the minimum wholesale cost of an item that can be purchased with a credit card?

- The length of a rectangle is twice the width of the rectangle. If the perimeter of the rectangle is no more than 15 mm, what is the greatest measure of the **length** of the rectangle?

- The sum of two consecutive integers is greater than the smaller integer increased by 10. What are the possible values for the **smaller** integer?

© 2003 BestQuest

Journal

1. Explain why the symbol “<” would not be used to model the following: John chose five less than y apples.
2. Write a problem that can be solved with the inequality: $0.25x < 500$.
3. Newt solved an inequality to answer a question about the maximum amount that a student would need to score on a test so that her average test score was not greater than 92. The answer to the equation was $s > 88$. Newt concluded that the student needed to score at least 88 points on her next test. Explain the error.
4. How would the answer to question 7 in the first section of this Independent Practice change if “no more than” was changed to “at least”?
5. Write an inequality that has no solution.

Cumulative Review

Simplify each expression.

- | | |
|---------------------------|----------------------------------------|
| 1. 5^2 _____ | 2. $\sqrt{121}$ _____ |
| 3. $\sqrt[3]{-125}$ _____ | 4. $(-4)^3$ _____ |
| 5. $\sqrt{-16}$ _____ | 6. 1^{87} _____ |
| 7. $(-10)^2$ _____ | 8. $-\sqrt{36}$ _____ |
| 9. k^0 _____ | 10. $\left(\frac{2}{3}\right)^3$ _____ |